



IFT122 gene

intraflagellar transport 122

Normal Function

The *IFT122* gene provides instructions for making a protein that is involved in the formation and maintenance of cilia, which are microscopic, finger-like projections that stick out from the surface of cells. Cilia participate in signaling pathways that transmit information within and between cells and are important for the development and function of many types of cells and tissues, including cells in the kidneys and liver and the light-sensitive tissue at the back of the eye (the retina). Cilia also play a role in the development of the bones, although the mechanism is not well understood.

The movement of substances within cilia and similar structures called flagella is known as intraflagellar transport. This process is essential for the assembly and maintenance of these cell structures. During intraflagellar transport, cells use molecules called IFT particles to carry materials to and from the tips of cilia. Each IFT particle is made up of two groups of IFT proteins: complex A and complex B. The protein produced from the *IFT122* gene forms part of IFT complex A (IFT-A). During intraflagellar transport, this complex carries materials from the tip to the base of cilia.

The IFT-A complex is essential for proper regulation of the Sonic Hedgehog signaling pathway, which is important for the growth and maturation (differentiation) of cells and the normal shaping (patterning) of many parts of the body, especially during embryonic development. The exact role of the complex in this pathway is unclear.

Health Conditions Related to Genetic Changes

cranioectodermal dysplasia

Mutations in the *IFT122* gene can cause cranioectodermal dysplasia. This condition is characterized by an elongated head (dolichocephaly) with a prominent forehead and other distinctive facial features; short bones; and abnormalities of certain tissues known as ectodermal tissues, which include the teeth, hair, nails, and skin.


At least six *IFT122* gene mutations have been found in people with cranioectodermal dysplasia. These mutations reduce the amount or function of the IFT122 protein. A shortage or reduction in activity of this component of the IFT-A complex impairs the function of the entire complex, disrupting transport of proteins and materials from the tips of cilia. As a result, assembly and maintenance of cilia is impaired, which leads to a smaller number of cilia and abnormalities in their shape and structure. Although the mechanism is unclear, a loss of normal cilia impedes proper development of bone and other tissues, leading to the features of cranioectodermal dysplasia. Some

researchers suggest that disrupted intraflagellar transport prevents signaling through the Sonic Hedgehog pathway, which could impact cell growth and other functions in several tissues throughout the body.

Chromosomal Location

Cytogenetic Location: 3q21.3-q22.1, which is the long (q) arm of chromosome 3 between positions 21.3 and 22.1

Molecular Location: base pairs 129,440,036 to 129,520,507 on chromosome 3 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- CED
- CED1
- IF122_HUMAN
- intraflagellar transport 122 homolog (Chlamydomonas)
- intraflagellar transport protein 122 homolog
- SPG
- WD repeat-containing protein 10
- WD repeat-containing protein 140
- WD repeat domain 10
- WDR10
- WDR10p
- WDR140

Additional Information & Resources

Educational Resources

- Molecular Cell Biology (fourth edition, 2000): Cilia and Flagella: Structure and Movement
<https://www.ncbi.nlm.nih.gov/books/NBK21698/>

GeneReviews

- Cranioectodermal Dysplasia
<https://www.ncbi.nlm.nih.gov/books/NBK154653>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28IFT122%5BTIAB%5D%29+OR+%28%28WDR10%5BTIAB%5D%29+OR+%28WDR10p%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- INTRAFLAGELLAR TRANSPORT 122, CHLAMYDOMONAS, HOMOLOG OF
<http://omim.org/entry/606045>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_IFT122.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=IFT122%5Bgene%5D>
- HGNC Gene Family: Intraflagellar transport proteins
<http://www.genenames.org/cgi-bin/genefamilies/set/615>
- HGNC Gene Family: WD repeat domain containing
<http://www.genenames.org/cgi-bin/genefamilies/set/362>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=13556
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/55764>
- UniProt
<http://www.uniprot.org/uniprot/Q9HBG6>

Sources for This Summary

- Arts H, Knoers N. Cranioectodermal Dysplasia. 2013 Sep 12. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJH, Bird TD, Fong CT, Mefford HC, Smith RJH, Stephens K, editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2016. Available from <http://www.ncbi.nlm.nih.gov/books/NBK154653/>
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24027799>
- OMIM: INTRAFLAGELLAR TRANSPORT 122, CHLAMYDOMONAS, HOMOLOG OF
<http://omim.org/entry/606045>
- Liem KF Jr, Ashe A, He M, Satir P, Moran J, Beier D, Wicking C, Anderson KV. The IFT-A complex regulates Shh signaling through cilia structure and membrane protein trafficking. *J Cell Biol.* 2012 Jun 11;197(6):789-800. doi: 10.1083/jcb.201110049.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22689656>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3373400/>
- Qin J, Lin Y, Norman RX, Ko HW, Eggenschwiler JT. Intraflagellar transport protein 122 antagonizes Sonic Hedgehog signaling and controls ciliary localization of pathway components. *Proc Natl Acad Sci U S A.* 2011 Jan 25;108(4):1456-61. doi: 10.1073/pnas.1011410108. Epub 2011 Jan 5.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21209331>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3029728/>
- Walczak-Sztulpa J, Eggenschwiler J, Osborn D, Brown DA, Emma F, Klingenberg C, Hennekam RC, Torre G, Garshasbi M, Tzschach A, Szczepanska M, Krawczynski M, Zachwieja J, Zwolinska D, Beales PL, Ropers HH, Latos-Bielenska A, Kuss AW. Cranioectodermal Dysplasia, Sensenbrenner syndrome, is a ciliopathy caused by mutations in the IFT122 gene. *Am J Hum Genet.* 2010 Jun 11;86(6):949-56. doi: 10.1016/j.ajhg.2010.04.012. Epub 2010 May 20.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20493458>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3032067/>

Reprinted from Genetics Home Reference:

<https://ghr.nlm.nih.gov/gene/IFT122>

Reviewed: November 2013

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services